

## 兔抗 BCL10 多克隆抗体

中文名称：兔抗 BCL10 多克隆抗体

英文名称： Anti-BCL10 rabbit polyclonal antibody

别名： CLAP; mE10; CIPER; c-E10; CARMEN

相关类别： 一抗

抗原： BCL10

储存： 冷冻（-20℃）

宿主： Rabbit

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

<b>Background:</b>	This gene was identified by its translocation in a case of mucosa-associated lymphoid tissue (MALT) lymphoma. The protein encoded by this gene contains a caspase recruitment domain (CARD), and has been shown to induce apoptosis and to activate NF- $\kappa$ B. This protein is reported to interact with other CARD domain containing proteins including CARD9, 10, 11 and 14, which are thought to function as upstream regulators in NF- $\kappa$ B signaling. This protein is found to form a complex with MALT1, a protein encoded by another gene known to be translocated in MALT lymphoma.
<b>Applications:</b>	ELISA, WB, IHC

<b>Name of antibody:</b>	BCL10
<b>Immunogen:</b>	Fusion protein of human BCL10
<b>Full name:</b>	B-cell CLL/lymphoma 10
<b>Synonyms :</b>	CLAP; mE10; CIPER; c-E10; CARMEN
<b>SwissProt:</b>	O95999
<b>ELISA Recommended dilution:</b>	2000-5000
<b>IHC positive control:</b>	Human thyroid cancer and human liver cancer
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	26 kDa
<b>WB Positive control:</b>	Hela cells
<b>WB Recommended dilution:</b>	500-2000



