

兔抗 CCL5 多克隆抗体

- 中文名称：兔抗 CCL5 多克隆抗体
- 英文名称：Anti-CCL5 rabbit polyclonal antibody
- 别名：C-C motif chemokine ligand 5 SISd; eoCP; SCYA5; RANTES; TCP228; D17S136E; SIS-delta
- 储存：冷冻（-20℃）
- 抗原：CCL5
- 宿主：Rabbit
- 相关类别：一抗
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:	<p>This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, functions as a chemoattractant for blood monocytes, memory T helper cells and eosinophils. It causes the release of histamine from basophils and activates eosinophils. This cytokine is one of the major HIV-suppressive factors produced by CD8+ cells. It functions as one of the natural ligands for the che</p>
--------------------	---

	mokine receptor chemokine (C-C motif) receptor 5 (CCR5), and it suppresses in vitro replication of the R5 strains of HIV-1, which use CCR5 as a coreceptor. Alternative splicing results in multiple transcript variants that encode different isoforms.
Applications:	ELISA, IHC
Name of antibody:	CCL5
Immunogen:	Synthetic peptide of human CCL5
Full name:	C-C motif chemokine ligand 5
Synonyms:	SISd; eoCP; SCYA5; RANTES; TCP228; D17S136E; SIS-delta
SwissProt:	P13501
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer;
IHC Recommend dilution:	25-100



